

REVERSE TRIANGLE INEQUALITY IN HILBERT C^* -MODULES

M. KHOSRAVI¹, H. MAHYAR² AND M. S. MOSLEHIAN³

ABSTRACT. We prove several versions of reverse triangle inequality in Hilbert C^* -modules. We show that if $e_1, \dots, e_m$ are vectors in a Hilbert module $\mathfrak{X}$ over a C^* -algebra $\mathfrak{A}$ with unit 1 such that $\langle e_i, e_j \rangle = 0$ ($1 \leq i \neq j \leq m$) and $\|e_i\| = 1$ ($1 \leq i \leq m$), and also $r_k, \rho_k \in \mathbb{R}$ ($1 \leq k \leq m$) and $x_1, \dots, x_n \in \mathfrak{X}$ satisfy

$$0 \leq r_k^2 \|x_j\| \leq \operatorname{Re} \langle r_k e_k, x_j \rangle, \quad 0 \leq \rho_k^2 \|x_j\| \leq \operatorname{Im} \langle \rho_k e_k, x_j \rangle,$$

then

$$\left[\sum_{k=1}^m (r_k^2 + \rho_k^2) \right]^{\frac{1}{2}} \sum_{j=1}^n \|x_j\| \leq \left\| \sum_{j=1}^n x_j \right\|,$$

and the equality holds if and only if

$$\sum_{j=1}^n x_j = \sum_{j=1}^n \|x_j\| \sum_{k=1}^m (r_k + i\rho_k) e_k.$$

1. INTRODUCTION AND PRELIMINARIES

The triangle inequality is one of the most fundamental inequalities in mathematics. Several mathematician have been investigated its generalizations and its reverses.

In 1917, Petrovitch [16] proved that for complex numbers $z_1, \dots, z_n$,

$$\left| \sum_{j=1}^n z_j \right| \geq \cos \theta \sum_{j=1}^n |z_j|,$$

where $0 < \theta < \frac{\pi}{2}$ and $\alpha - \theta < \arg z_j < \alpha + \theta$ ($1 \leq j \leq n$) for a given real number α .

The first generalization of the reverse triangle inequality in Hilbert spaces was given by Diaz and Matcalf [5]. They proved that for $x_1, \dots, x_n$ in a Hilbert space H , if e is a unit vector

2000 *Mathematics Subject Classification.* Primary 46L08; Secondary 15A39, 26D15, 46L05, 51M16.

Key words and phrases. Triangle inequality, Reverse inequality, Hilbert C^* -module, C^* -algebra.

of H such that $0 \leq r \leq \frac{\operatorname{Re}\langle x_j, e \rangle}{\|x_j\|}$ for some $r \in \mathbb{R}$ and each $1 \leq j \leq n$, then

$$r \sum_{j=1}^n \|x_j\| \leq \left\| \sum_{j=1}^n x_j \right\|.$$

Moreover, the equality holds if and only if $\sum_{j=1}^n x_j = r \sum_{j=1}^n \|x_j\| e$.

Recently, a number of mathematicians have represented several refinements of the reverse triangle inequality in Hilbert spaces and normed spaces. See [1, 2, 4, 7, 8, 10, 13, 15]. Recently A discussion of C^* -valued triangle inequality in Hilbert C^* -modules was given in [3]. Our aim is to give some generalizations of results of Dragomir in Hilbert spaces to the framework of Hilbert C^* -modules. For this purpose, we first recall some fundamental definitions in the theory of Hilbert C^* -modules.

Suppose that $\mathfrak{A}$ is a C^* -algebra and $\mathfrak{X}$ is a linear space, which is an algebraic right $\mathfrak{A}$ -module. The space $\mathfrak{X}$ is called a pre-Hilbert $\mathfrak{A}$ -module (or an inner product $\mathfrak{A}$ -module) if there exists an $\mathfrak{A}$ -valued inner product $\langle \cdot, \cdot \rangle : \mathfrak{X} \times \mathfrak{X} \rightarrow \mathfrak{A}$ with the following properties:

- (i) $\langle x, x \rangle \geq 0$ and $\langle x, x \rangle = 0$ if and only if $x = 0$
- (ii) $\langle x, \lambda y + z \rangle = \lambda \langle x, y \rangle + \langle x, z \rangle$
- (iii) $\langle x, ya \rangle = \langle x, y \rangle a$
- (iv) $\langle x, y \rangle^* = \langle y, x \rangle$

for all $x, y, z \in \mathfrak{X}$, $a \in \mathfrak{A}$, $\lambda \in \mathbb{C}$. By (ii) and (iv), $\langle \cdot, \cdot \rangle$ is conjugate linear in the first variable. Using the Cauchy–Schwartz inequality $\langle y, x \rangle \langle x, y \rangle \leq \|\langle x, x \rangle\| \|\langle y, y \rangle\|$ [11, Page 5], it follows that $\|x\| = \|\langle x, x \rangle\|^{\frac{1}{2}}$ is a norm on $\mathfrak{X}$ making it into a right normed module. The pre-Hilbert module $\mathfrak{X}$ is called a Hilbert $\mathfrak{A}$ -module if it is complete with respect to this norm. Notice that the inner structure of a C^* -algebra is essentially more complicated than complex numbers. For instance, the notations such as orthogonality and theorems such as Riesz' representation in the complex Hilbert space theory cannot simply be generalized or transferred to the theory of Hilbert C^* -modules.

One may define an “ $\mathfrak{A}$ -valued norm” $|\cdot|$ by $|x| = \langle x, x \rangle^{1/2}$. Clearly, $\| |x| \| = \|x\|$ for each $x \in \mathfrak{X}$. It is known that $|\cdot|$ does not satisfy the triangle inequality in general. See [11, 12] for more information on Hilbert C^* -modules.

We also use the elementary C^* -algebra theory, in particular we utilize this property that if $a \leq b$ then $a^{1/2} \leq b^{1/2}$, where a, b are positive elements of a C^* -algebra $\mathfrak{A}$. We also repeatedly apply the following known relation:

$$\frac{1}{2}(aa^* + a^*a) = (\operatorname{Re} a)^2 + (\operatorname{Im} a)^2, \quad (\diamond)$$

where a is an arbitrary element of $\mathfrak{A}$. For details on C^* -algebra theory we referred the readers to [14].

Throughout this paper, we assume that $\mathfrak{A}$ is a unital C^* -algebra with unit 1 and for every $\lambda \in \mathbb{C}$, we write $\lambda 1$ as λ .

2. MULTIPLICATIVE REVERSE OF THE TRIANGLE INEQUALITY

Utilizing some C^* -algebraic techniques we present our first result as a generalization of [7, Theorem 2.3].

Theorem 2.1. *Let $\mathfrak{A}$ be a C^* -algebra, let $\mathfrak{X}$ be a Hilbert $\mathfrak{A}$ -module and let $x_1, \dots, x_n \in \mathfrak{X}$. If there exist real numbers $k_1, k_2 \geq 0$ with*

$$0 \leq k_1 \|x_j\| \leq \operatorname{Re} \langle e, x_j \rangle, \quad 0 \leq k_2 \|x_j\| \leq \operatorname{Im} \langle e, x_j \rangle,$$

for some $e \in \mathfrak{X}$ with $|e| \leq 1$ and all $1 \leq j \leq n$, then

$$(k_1^2 + k_2^2)^{\frac{1}{2}} \sum_{j=1}^n \|x_j\| \leq \left\| \sum_{j=1}^n x_j \right\|. \quad (2.1)$$

Proof. Applying the Cauchy–Schwarz inequality, we get

$$|\langle e, \sum_{j=1}^n x_j \rangle|^2 \leq \|e\|^2 \left\| \sum_{j=1}^n x_j \right\|^2 \leq \left\| \sum_{j=1}^n x_j \right\|^2,$$

and

$$|\langle \sum_{j=1}^n x_j, e \rangle|^2 \leq \left\| \sum_{j=1}^n x_j \right\|^2 |e|^2 \leq \left\| \sum_{j=1}^n x_j \right\|^2,$$

whence

$$\begin{aligned}
\left\| \sum_{j=1}^n x_j \right\|^2 &\geq \frac{1}{2} \left(|\langle e, \sum_{j=1}^n x_j \rangle|^2 + |\langle \sum_{j=1}^n x_j, e \rangle|^2 \right) \\
&= \frac{1}{2} \left(\langle e, \sum_{j=1}^n x_j \rangle^* \langle e, \sum_{j=1}^n x_j \rangle + \langle \sum_{j=1}^n x_j, e \rangle^* \langle \sum_{j=1}^n x_j, e \rangle \right) \\
&= (\operatorname{Re} \langle e, \sum_{j=1}^n x_j \rangle)^2 + (\operatorname{Im} \langle e, \sum_{j=1}^n x_j \rangle)^2 \quad (\text{by } (\diamond)) \\
&= (\operatorname{Re} \sum_{j=1}^n \langle e, x_j \rangle)^2 + (\operatorname{Im} \sum_{j=1}^n \langle e, x_j \rangle)^2 \\
&\geq k_1^2 (\sum_{j=1}^n \|x_j\|)^2 + k_2^2 (\sum_{j=1}^n \|x_j\|)^2 \\
&= (k_1^2 + k_2^2) (\sum_{j=1}^n \|x_j\|)^2.
\end{aligned}$$

□

Using the same argument as in the proof of Theorem 2.1 one can obtain the following result, where k_1, k_2 are hermitian elements of $\mathfrak{A}$.

Theorem 2.2. *If the vectors $x_1, \dots, x_n \in \mathfrak{X}$ satisfy the conditions*

$$0 \leq k_1^2 \|x_j\|^2 \leq (\operatorname{Re} \langle e, x_j \rangle)^2, \quad 0 \leq k_2^2 \|x_j\|^2 \leq (\operatorname{Im} \langle e, x_j \rangle)^2,$$

for some hermitian elements k_1, k_2 in $\mathfrak{A}$, some $e \in \mathfrak{X}$ with $|e| \leq 1$ and all $1 \leq j \leq n$ then the inequality 2.1 holds.

One may observe an integral version of inequality (2.1) as follows:

Corollary 2.3. *Suppose that $\mathfrak{X}$ is a Hilbert $\mathfrak{A}$ -module and $f : [a, b] \rightarrow \mathfrak{X}$ is strongly measurable such that the Lebesgue integral $\int_a^b \|f(t)\| dt$ exists and is finite. If there exist self-adjoint elements a_1, a_2 in $\mathfrak{A}$ with*

$$a_1^2 \|f(t)\|^2 \leq \operatorname{Re} \langle f(t), e \rangle^2, \quad a_2^2 \|f(t)\|^2 \leq \operatorname{Im} \langle f(t), e \rangle^2 \quad (\text{a.e. } t \in [a, b]),$$

where $e \in \mathfrak{X}$ with $|e| \leq 1$, then

$$(a_1^2 + a_2^2)^{\frac{1}{2}} \int_a^b \|f(t)\| dt \leq \left\| \int_a^b f(t) dt \right\|.$$

Now we prove a useful lemma, which is frequently applied in the next theorems (see also [3]).

Lemma 2.4. *Let $\mathfrak{X}$ be a Hilbert $\mathfrak{A}$ -module and let $x, y \in \mathfrak{X}$. If $|\langle x, y \rangle| = \|x\| \|y\|$, then*

$$y = \frac{x \langle x, y \rangle}{\|x\|^2}.$$

Proof. For $x, y \in \mathfrak{X}$ we have

$$\begin{aligned} 0 &\leq \left| y - \frac{x \langle x, y \rangle}{\|x\|^2} \right|^2 = \left\langle y - \frac{x \langle x, y \rangle}{\|x\|^2}, y - \frac{x \langle x, y \rangle}{\|x\|^2} \right\rangle \\ &= \langle y, y \rangle - \frac{1}{\|x\|^2} \langle y, x \rangle \langle x, y \rangle + \frac{1}{\|x\|^4} \langle y, x \rangle \langle x, x \rangle \langle x, y \rangle - \frac{1}{\|x\|^2} \langle y, x \rangle \langle x, y \rangle \\ &\leq |y|^2 - \frac{1}{\|x\|^2} |\langle x, y \rangle|^2 = |y|^2 - \frac{1}{\|x\|^2} \|x\|^2 \|y\|^2 \\ &= |y|^2 - \|y\|^2 \leq 0, \end{aligned}$$

whence $\left| y - \frac{x \langle x, y \rangle}{\|x\|^2} \right| = 0$. Hence $y = \frac{x \langle x, y \rangle}{\|x\|^2}$. □

Using the Cauchy–Schwarz inequality, we have the following theorem for Hilbert modules, which is similar to [1, Theorem 2.5].

Theorem 2.5. *Let $e_1, \dots, e_m$ be a family of vectors in a Hilbert module $\mathfrak{X}$ over a C^* -algebra $\mathfrak{A}$ such that $\langle e_i, e_j \rangle = 0$ ($1 \leq i \neq j \leq m$) and $\|e_i\| = 1$ ($1 \leq i \leq m$). Suppose that $r_k, \rho_k \in \mathbb{R}$ ($1 \leq k \leq m$) and that the vectors $x_1, \dots, x_n \in \mathfrak{X}$ satisfy*

$$0 \leq r_k^2 \|x_j\| \leq \operatorname{Re} \langle r_k e_k, x_j \rangle, \quad 0 \leq \rho_k^2 \|x_j\| \leq \operatorname{Im} \langle \rho_k e_k, x_j \rangle,$$

Then

$$\left[\sum_{k=1}^m (r_k^2 + \rho_k^2) \right]^{\frac{1}{2}} \sum_{j=1}^n \|x_j\| \leq \left\| \sum_{j=1}^n x_j \right\|, \quad (2.2)$$

and the equality holds if and only if

$$\sum_{j=1}^n x_j = \sum_{j=1}^n \|x_j\| \sum_{k=1}^m (r_k + i \rho_k) e_k. \quad (2.3)$$

Proof. There is nothing to prove if $\sum_{k=1}^m (r_k^2 + \rho_k^2) = 0$. Assume that $\sum_{k=1}^m (r_k^2 + \rho_k^2) \neq 0$. From the hypothesis we have

$$\begin{aligned}
\left(\sum_{k=1}^m (r_k^2 + \rho_k^2) \right)^2 \left(\sum_{j=1}^n \|x_j\| \right)^2 &\leq \left(\operatorname{Re} \langle \sum_{k=1}^m r_k e_k, \sum_{j=1}^n x_j \rangle + \operatorname{Im} \langle \sum_{k=1}^m \rho_k e_k, \sum_{j=1}^n x_j \rangle \right)^2 \\
&= \left(\operatorname{Re} \langle \sum_{j=1}^n x_j, \sum_{k=1}^m (r_k + i\rho_k) e_k \rangle \right)^2 \\
&\quad \left(\text{by } \operatorname{Im}(a) = \operatorname{Re}(ia^*), \operatorname{Re}(a^*) = \operatorname{Re}(a) \quad (a \in \mathfrak{A}) \right) \\
&\leq \frac{1}{2} \left| \langle \sum_{j=1}^n x_j, \sum_{k=1}^m (r_k + i\rho_k) e_k \rangle \right|^2 \\
&\quad + \left| \langle \sum_{k=1}^m (r_k + i\rho_k) e_k, \sum_{j=1}^n x_j \rangle \right|^2 \quad (\text{by } (\diamond)) \\
&\leq \frac{1}{2} \left\| \sum_{j=1}^n x_j \right\|^2 \left\| \sum_{k=1}^m (r_k + i\rho_k) e_k \right\|^2 \\
&\quad + \left\| \sum_{k=1}^m (r_k + i\rho_k) e_k \right\|^2 \left\| \sum_{j=1}^n x_j \right\|^2 \\
&\leq \left\| \sum_{j=1}^n x_j \right\|^2 \left\| \sum_{k=1}^m (r_k + i\rho_k) e_k \right\|^2 \\
&\quad (\text{by } |a| \leq \|a\| \quad (a \in \mathfrak{A})) \\
&= \left\| \sum_{j=1}^n x_j \right\|^2 \left\| \langle \sum_{k=1}^m (r_k + i\rho_k) e_k, \sum_{k=1}^m (r_k + i\rho_k) e_k \rangle \right\| \\
&= \left\| \sum_{j=1}^n x_j \right\|^2 \sum_{k=1}^m |r_k + i\rho_k|^2 \|e_k\|^2 \\
&= \left\| \sum_{j=1}^n x_j \right\|^2 \sum_{k=1}^m (r_k^2 + \rho_k^2).
\end{aligned}$$

Hence

$$\left[\sum_{k=1}^m (r_k^2 + \rho_k^2) \right] \left(\sum_{j=1}^n \|x_j\| \right)^2 \leq \left\| \sum_{j=1}^n x_j \right\|^2.$$

By taking square roots the desired result follows.

Clearly we have equality in 2.2 if condition 2.3 holds. To see the converse, first note that if equality holds in 2.2, then all inequalities in the relations above should be equality. Therefore

$$r_k^2 \|x_j\| = \operatorname{Re} \langle r_k e_k, x_j \rangle, \quad \rho_k^2 \|x_j\| = \operatorname{Im} \langle \rho_k e_k, x_j \rangle,$$

$$\operatorname{Re} \left\langle \sum_{j=1}^n x_j, \sum_{k=1}^m (r_k + i\rho_k) e_k \right\rangle = \left\langle \sum_{j=1}^n x_j, \sum_{k=1}^m (r_k + i\rho_k) e_k \right\rangle,$$

and also

$$\left| \left\langle \sum_{k=1}^m (r_k + i\rho_k) e_k, \sum_{j=1}^n x_j \right\rangle \right| = \left\| \sum_{j=1}^n x_j \right\| \left\| \sum_{k=1}^m (r_k + i\rho_k) e_k \right\|.$$

From Lemma 2.4 and these equalities we have

$$\begin{aligned}
 \sum_{j=1}^n x_j &= \frac{\sum_{k=1}^m (r_k + i\rho_k)e_k}{\|\sum_{k=1}^m (r_k + i\rho_k)e_k\|^2} \langle \sum_{k=1}^m (r_k + i\rho_k)e_k, \sum_{j=1}^n x_j \rangle \\
 &= \frac{\sum_{k=1}^m (r_k + i\rho_k)e_k}{\sum_{k=1}^m (r_k^2 + \rho_k^2)} \operatorname{Re} \langle \sum_{k=1}^m (r_k + i\rho_k)e_k, \sum_{j=1}^n x_j \rangle \\
 &= \frac{\sum_{k=1}^m (r_k + i\rho_k)e_k}{\sum_{k=1}^m (r_k^2 + \rho_k^2)} \sum_{k=1}^m \sum_{j=1}^n (r_k^2 \|x_j\| + \rho_k^2 \|x_j\|) \\
 &= \sum_{j=1}^n \|x_j\| \sum_{k=1}^m (r_k + i\rho_k)e_k,
 \end{aligned}$$

which is the desired result. $\square$

In the next results of this section, we assume that $\mathfrak{X}$ is a right Hilbert $\mathfrak{A}$ -module, which is an algebraic left A -module subject to

$$\langle x, ay \rangle = a \langle x, y \rangle \quad (x, y \in \mathfrak{X}, a \in \mathfrak{A}). \quad (\dagger)$$

For example if $\mathfrak{A}$ is a unital C^* -algebra and $\mathfrak{I}$ is a commutative right ideal of $\mathfrak{A}$, then $\mathfrak{I}$ is a right Hilbert module over $\mathfrak{A}$ and

$$\langle x, ay \rangle = x^*(ay) = ax^*y = a \langle x, y \rangle \quad (x, y \in \mathfrak{I}, a \in \mathfrak{A}).$$

The next theorem is a refinement of [7, Theorem 2.1]. To prove it we need the following lemma.

Lemma 2.6. *Let $\mathfrak{X}$ be a Hilbert $\mathfrak{A}$ -module and $e_1, \dots, e_n \in \mathfrak{X}$ be a family of vectors such that $\langle e_i, e_j \rangle = 0$ ($i \neq j$) and $\|e_i\| = 1$. If $x \in \mathfrak{X}$, then*

$$|x|^2 \geq \sum_{k=1}^n |\langle e_k, x \rangle|^2 \quad \text{and} \quad |x|^2 \geq \sum_{k=1}^n |\langle x, e_k \rangle|^2.$$

Proof. The first result follows from the following inequality:

$$\begin{aligned}
 0 \leq |x - \sum_{k=1}^n e_k \langle e_k, x \rangle|^2 &= \langle x - \sum_{k=1}^n e_k \langle e_k, x \rangle, x - \sum_{j=1}^n e_j \langle e_j, x \rangle \rangle \\
 &= \langle x, x \rangle + \sum_{k=1}^n \sum_{j=1}^n \langle e_k, x \rangle^* \langle e_k, e_j \rangle \langle e_j, x \rangle - 2 \sum_{k=1}^n |\langle e_k, x \rangle|^2 \\
 &= \langle x, x \rangle + \sum_{k=1}^n \langle e_k, x \rangle^* \langle e_k, e_k \rangle \langle e_k, x \rangle - 2 \sum_{k=1}^n |\langle e_k, x \rangle|^2 \\
 &\leq |x|^2 + \sum_{k=1}^n \langle e_k, x \rangle^* \langle e_k, x \rangle - 2 \sum_{k=1}^n |\langle e_k, x \rangle|^2 \\
 &= |x|^2 - \sum_{k=1}^n |\langle e_k, x \rangle|^2.
 \end{aligned}$$

By considering $|x - \sum_{k=1}^n \langle e_k, x \rangle e_k|^2$, similarly, we get the second one. $\square$

Now we are able to prove the next theorem without using the Cauchy–Schwarz inequality.

Theorem 2.7. *Let $e_1, \dots, e_m \in \mathfrak{X}$ be a family of vectors with $\langle e_i, e_j \rangle = 0$ ($1 \leq i \neq j \leq m$) and $\|e_i\| = 1$ ($1 \leq i \leq m$). If the vectors $x_1, \dots, x_n \in \mathfrak{X}$ satisfy the conditions*

$$0 \leq r_k \|x_j\| \leq \operatorname{Re} \langle e_k, x_j \rangle, \quad 0 \leq \rho_k \|x_j\| \leq \operatorname{Im} \langle e_k, x_j \rangle \quad (1 \leq j \leq n, 1 \leq k \leq m), \quad (2.4)$$

where $r_k, \rho_k \in [0, \infty)$ ($1 \leq k \leq m$), then

$$\left[\sum_{k=1}^m (r_k^2 + \rho_k^2) \right]^{\frac{1}{2}} \sum_{j=1}^n \|x_j\| \leq \left| \sum_{j=1}^n x_j \right|. \quad (2.5)$$

Proof. Applying the previous lemma for $x = \sum_{j=1}^n x_j$, we obtain

$$\begin{aligned} \left| \sum_{j=1}^n x_j \right|^2 &\geq \frac{1}{2} \left(\sum_{k=1}^m |\langle e_k, \sum_{j=1}^n x_j \rangle|^2 + \sum_{k=1}^m |\langle \sum_{j=1}^n x_j, e_k \rangle|^2 \right) \\ &= \sum_{k=1}^m \frac{1}{2} \left(\langle e_k, \sum_{j=1}^n x_j \rangle^* \langle e_k, \sum_{j=1}^n x_j \rangle + \langle \sum_{j=1}^n x_j, e_k \rangle^* \langle \sum_{j=1}^n x_j, e_k \rangle \right) \\ &= \sum_{k=1}^m (\operatorname{Re} \langle e_k, \sum_{j=1}^n x_j \rangle)^2 + (\operatorname{Im} \langle e_k, \sum_{j=1}^n x_j \rangle)^2 \quad (\text{by } (\diamond)) \\ &= \sum_{k=1}^m (\operatorname{Re} \sum_{j=1}^n \langle e_k, x_j \rangle)^2 + (\operatorname{Im} \sum_{j=1}^n \langle e_k, x_j \rangle)^2 \\ &\geq \sum_{k=1}^m (r_k^2 (\sum_{j=1}^n \|x_j\|)^2 + \rho_k^2 (\sum_{j=1}^n \|x_j\|)^2) \quad (\text{by (2.4)}) \\ &= \sum_{k=1}^m (r_k^2 + \rho_k^2) (\sum_{j=1}^n \|x_j\|)^2. \end{aligned}$$

□

Proposition 2.8. *In Theorem 2.7, if $\langle e_k, e_k \rangle = 1$, then the equality holds in (2.5) if and only if*

$$\sum_{j=1}^n x_j = \left(\sum_{j=1}^n \|x_j\| \right) \sum_{k=1}^m (r_k + i\rho_k) e_k. \quad (2.6)$$

Proof. If (2.6) holds, then inequality in (2.5) turns trivially into equality.

Next, assume that equality holds in (2.5). Then two inequalities in the proof of Theorem 2.7 should be equality. Hence

$$\left| \sum_{j=1}^n x_j \right|^2 = \sum_{k=1}^m \left| \langle e_k, \sum_{j=1}^n x_j \rangle \right|^2 \quad \text{and} \quad \left| \sum_{j=1}^n x_j \right|^2 = \sum_{k=1}^m \left| \langle \sum_{j=1}^n x_j, e_k \rangle \right|^2,$$

which is equivalent to

$$\sum_{j=1}^n x_j = \sum_{k=1}^m \sum_{j=1}^n e_k \langle e_k, x_j \rangle = \sum_{k=1}^m \sum_{j=1}^n \langle e_k, x_j \rangle e_k,$$

and also

$$r_k \|x_j\| = \operatorname{Re}\langle e_k, x_j \rangle, \quad \rho_k \|x_j\| = \operatorname{Im}\langle e_k, x_j \rangle.$$

So

$$\sum_{j=1}^n x_j = \sum_{k=1}^m \sum_{j=1}^n e_k \langle e_k, x_j \rangle = \sum_{k=1}^m \sum_{j=1}^n e_k (r_k + i\rho_k) \|x_j\| = \left(\sum_{j=1}^n \|x_j\| \right) \sum_{k=1}^m (r_k + i\rho_k) e_k.$$

□

3. ADDITIVE REVERSE OF THE TRIANGLE INEQUALITY

There are some versions of additive reverse of triangle inequality. In [6], Dragomir established the following theorem:

Theorem 3.1. *Let $\{e_k\}_{k=1}^m$ be a family of orthonormal vectors in Hilbert space H and $M_{jk} \geq 0$ ($1 \leq j \leq n, 1 \leq k \leq m$) such that*

$$\|x_j\| - \operatorname{Re}\langle e_k, x_j \rangle \leq M_{jk},$$

for each $1 \leq j \leq n$ and $1 \leq k \leq m$. Then

$$\sum_{j=1}^n \|x_j\| \leq \frac{1}{\sqrt{m}} \left\| \sum_{j=1}^n x_j \right\| + \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk};$$

and the equality holds if and only if

$$\sum_{j=1}^n \|x_j\| \geq \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk},$$

and

$$\sum_{j=1}^n x_j = \left(\sum_{j=1}^n \|x_j\| - \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk} \right) \sum_{k=1}^m e_k.$$

We can prove this theorem for Hilbert C^* -modules with some different techniques.

Theorem 3.2. *Let $\{e_k\}_{k=1}^m$ be a family of vectors in Hilbert module $\mathfrak{X}$ over a C^* -algebra $\mathfrak{A}$ with unit 1 with $|e_k| \leq 1$ ($1 \leq k \leq m$) and $\langle e_i, e_j \rangle = 0$ ($1 \leq i \neq j \leq m$) and $x_j \in \mathfrak{X}$ ($1 \leq j \leq n$). If for some scalars $M_{jk} \geq 0$ ($1 \leq j \leq n, 1 \leq k \leq m$),*

$$\|x_j\| - \operatorname{Re}\langle e_k, x_j \rangle \leq M_{jk} \quad (1 \leq j \leq n, 1 \leq k \leq m), \quad (3.1)$$

then

$$\sum_{j=1}^n \|x_j\| \leq \frac{1}{\sqrt{m}} \left\| \sum_{j=1}^n x_j \right\| + \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk}. \quad (3.2)$$

Moreover, if $|e_k| = 1$ ($1 \leq k \leq m$), then the equality in (3.2) holds if and only if

$$\sum_{j=1}^n \|x_j\| \geq \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk}, \quad (3.3)$$

and

$$\sum_{j=1}^n x_j = \left(\sum_{j=1}^n \|x_j\| - \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk} \right) \sum_{k=1}^m e_k. \quad (3.4)$$

Proof. Taking the summation in (3.1) over j from 1 to n , we obtain

$$\sum_{j=1}^n \|x_j\| \leq \operatorname{Re} \langle e_k, \sum_{j=1}^n x_j \rangle + \sum_{j=1}^n M_{jk},$$

for each $k \in \{1, \dots, m\}$. Summing these inequalities over k from 1 to m , we deduce

$$\sum_{j=1}^n \|x_j\| \leq \frac{1}{m} \operatorname{Re} \langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \rangle + \frac{1}{m} \sum_{k=1}^m \sum_{j=1}^n M_{jk}. \quad (3.5)$$

Using the Cauchy–Schwarz we obtain

$$\begin{aligned} \left(\operatorname{Re} \langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \rangle \right)^2 &\leq \frac{1}{2} (|\langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \rangle|^2 + |\langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \rangle^*|^2) \quad (\text{by } (\diamond)) \\ &\leq \frac{1}{2} (\| \sum_{k=1}^m e_k \|^2 \| \sum_{j=1}^n x_j \|^2 + \| \sum_{k=1}^m e_k \|^2 \| \sum_{j=1}^n x_j \|^2) \\ &\leq \| \sum_{k=1}^m e_k \|^2 \| \sum_{j=1}^n x_j \|^2 \\ &\leq m \| \sum_{j=1}^n x_j \|^2, \end{aligned} \quad (3.6)$$

since

$$\left\| \sum_{k=1}^m e_k \right\|^2 = \left\| \langle \sum_{k=1}^m e_k, \sum_{k=1}^m e_k \rangle \right\| = \left\| \sum_{k=1}^m \sum_{l=1}^m \langle e_k, e_l \rangle \right\| = \left\| \sum_{k=1}^m |e_k|^2 \right\| \leq m.$$

Using (3.6) in (3.5), we deduce the desired inequality.

If (3.3) and (3.4) hold, then

$$\begin{aligned} \frac{1}{\sqrt{m}} \left\| \sum_{j=1}^n x_j \right\| &= \frac{1}{\sqrt{m}} \left(\sum_{j=1}^n \|x_j\| - \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk} \right) \left\| \sum_{k=1}^m e_k \right\| \\ &= \sum_{j=1}^n \|x_j\| - \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk}, \end{aligned}$$

and then the equality in 3.2 holds true.

Conversely, if the equality holds in (3.2), then obviously (3.3) is valid and we have equalities all over in the proof above. This means that

$$\begin{aligned} \|x_j\| - \operatorname{Re}\langle e_k, x_j \rangle &= M_{jk}, \\ \operatorname{Re}\langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \rangle &= \langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \rangle, \end{aligned}$$

and

$$|\langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \rangle| = \|\sum_{k=1}^m e_k\| \|\sum_{j=1}^n x_j\|.$$

It follows from Lemma 2.4 and the previous relations that

$$\begin{aligned} \sum_{j=1}^n x_j &= \frac{\sum_{k=1}^m e_k}{\|\sum_{k=1}^m e_k\|^2} \langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \rangle \\ &= \frac{\sum_{k=1}^m e_k}{m} \operatorname{Re}\langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \rangle \\ &= \frac{\sum_{k=1}^m e_k}{m} \sum_{k=1}^m \sum_{j=1}^n (\|x_j\| - M_{jk}) \\ &= \left(\sum_{j=1}^n \|x_j\| - \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk} \right) \sum_{k=1}^m e_k. \end{aligned}$$

□

REFERENCES

- [1] A.H. Ansari and M.S. Moslehian, *Refinements of reverse triangle inequality in inner product spaces*, J. Inequal. Pure Appl. Math., **6**(3) (2005), Article 64.
- [2] A.H. Ansari and M.S. Moslehian, *More on reverse triangle inequality in inner product spaces*, Inter J. Math. Math Sci., **18** (2005), 2883–2893.
- [3] Lj. Arambačić and R. Rajić, *On the C^* -valued triangle equality and inequality in Hilbert C^* -modules*, Acta Math. Hungar. **119** (2008), no. 4, 373–380.
- [4] I. Brnetić, S. Dragomir, R. Hoxha and J. Pečarić, *A reverse of the triangle inequality in inner product spaces and applications for polynomials*, Aust. J. Math. Anal. Appl., **3**(2) (2006), Art. 9.
- [5] J.B. Diaz and F.T. Metcalf, *A complementary triangle inequality in Hilbert and Banach spaces*, Proc. Amer. Math. Soc., **17**(1) (1966), 88–97.
- [6] S.S. Dragomir, *Reverses of the triangle inequality in inner product spaces*, Linear Algebra Appl., **402** (2005), 245–254.
- [7] S.S. Dragomir, *Some reverses of the generalized triangle inequality in complex inner product spaces*, RGMIA Res. Rep. Coll., **7** (2004), supplement, Article 7.

- [8] S.S. Dragomir, *Reverse of the continuous triangle inequality for Bochner integral in complex Hilbert spaces*, J. Math. Anal. Appl., **329** (2007), 65–76.
- [9] J. Karamata, *Teorija i Praksa Stieltjesova Integrala (Sebro-Coratian)(Stieltjes Integral, Theory and Practice)*, SANU, Posebna izdanja, **154**, Beograd, 1949.
- [10] M. Kato, K.S. Saito and T. Tamura, *Sharp triangle inequality and its reverse in Banach spaces*, Math. Inequal. Appl., **10** (2007), 451–460.
- [11] E.C. Lance, *Hilbert C^* -modules*, London Mathematical Society Lecture Note Series, 210, Cambridge University Press, Cambridge, 1995.
- [12] V.M. Manuilov and E.V. Troitsky, *Hilbert C^* -modules*, Translations of Mathematical Monographs, 226. American Mathematical Society, Providence, RI, 2005.
- [13] D.S. Mitrinović, J. E. Pečarić and A. M. Fink, *Classical and new inequalities in analysis*, Kluwer Academic, Dordrecht, 1993.
- [14] J.G. Murphy, *C^* -Algebras and Operator Theory*, Academic Press, Boston, 1990.
- [15] M. Nakai and T. Tada, *The reverse triangle inequality in normed spaces*, New Zealand J. Math. **25** (1996), no. 2, 181–193.
- [16] M. Petrovich, *Module d'une somme*, L'Enseignement Math., **19** (1917), 53–56.

MARYAM KHOSRAVI: DEPARTMENT OF MATHEMATICS, TEACHER TRAINING UNIVERSITY, TAHRAN, IRAN.

E-mail address: khosravi_m@saba.tmu.ac.ir

HAKIMEH MAHYAR: DEPARTMENT OF MATHEMATICS, TEACHER TRAINING UNIVERSITY, TAHRAN, IRAN.

E-mail address: mahyar@saba.tmu.ac.ir

MOHAMMAD SAL MOSLEHIAN: DEPARTMENT OF PURE MATHEMATICS AND CENTRE OF EXCELLENCE IN ANALYSIS ON ALGEBRAIC STRUCTURES (CEAAS), FERDOWSI UNIVERSITY OF MASHHAD, P. O. BOX 1159, MASHHAD 91775, IRAN.

E-mail address: moslehian@ferdowsi.um.ac.ir and moslehian@ams.org